

Week 5 Practice

CSCI 567 Machine Learning

Spring 2025

Instructor: Haipeng Luo

1. MULTIPLE-CHOICE QUESTIONS: One or more correct choice(s) for each question.

1.1. Which of the following is/are **true** about neural nets?

- (A) A neural net with one hidden layer and a fixed number of neurons can represent any continuous function.
- (B) A fully connected feedforward neural net without nonlinear activation functions is the same as a linear model.
- (C) Dropping random neurons in each iteration of Backpropagation helps prevent overfitting.
- (D) A max-pooling layer with a 2×2 filter has 4 parameters to be learned.

1.2. Which of the following can help prevent overfitting in neural nets?

- (A) Retraining on the same data many times.
- (B) Using a validation set for early stopping.
- (C) Data augmentation.
- (D) Training until you get the smallest training error.

1.3. Suppose a convolution layer takes a 8×8 image with 3 channels as input and outputs a $4 \times 4 \times 8$ volume. Which of the following is a possible configuration of this layer?

- (A) One 4×4 filter with depth 8, stride 2, 1 pixel of zero-padding.
- (B) Three 4×4 filters with depth 8, stride 2, no zero-padding.
- (C) Eight 4×4 filters with depth 3, stride 2, 1 pixel of zero-padding.
- (D) Eight 4×4 filters with depth 3, stride 1, 1 pixel of zero-padding.

- 1.4. How many parameters do we need to learn for the following network structure? An $32 \times 32 \times 3$ image input, followed by a convolution layer with 3 filters of size 3×3 (stride 1, 1 pixel of zero-padding), then another convolution layer with 4 filters of size 2×2 (stride 2, no zero-padding), and finally a max-pooling layer with a 2×2 filter (stride 1, no zero-padding). (Note: the depth of all filters are not explicitly spelled out, and we assume no bias/intercept terms are used.)
- (A) 43
 (B) 97
 (C) 129
 (D) 145

Ans: C. $3 \times (3 \times 3 \times 3) + 4 \times (2 \times 2 \times 3) = 129$.

- 1.5. What is the final output dimension of the last question?
- (A) $15 \times 15 \times 1$
 (B) $16 \times 16 \times 4$
 (C) $32 \times 32 \times 1$
 (D) $15 \times 15 \times 4$

- 1.6. Which of the following is/are **true** about kernel functions?

- (A) If k_1 and k_2 are valid kernels, then so is $c_1 k_1 + c_2 k_2$ for any $c_1, c_2 \geq 0$.
 (B) Kernel functions must be symmetric, i.e., $k(\mathbf{x}, \mathbf{x}') = k(\mathbf{x}', \mathbf{x})$.
 (C) If k is a kernel, then $-k$ is a kernel too.
 (D) If k is a kernel, then $\ln(k)$ is a kernel too.

- 1.7. Which of the following are not kernel functions?

- (A) $k(x, x') = (\mathbf{x}^T \mathbf{x}')^2$
 (B) $k(x, x') = -\|\mathbf{x} - \mathbf{x}'\|_2^2$
 (C) $k(x, x') = (\mathbf{x}^T \mathbf{x}' + 1)^2$
 (D) $k(x, x') = \|\mathbf{x} - \mathbf{x}'\|_2^2$

- 1.8. Vovk's real polynomial kernel $k : \mathbb{R}^D \times \mathbb{R}^D \rightarrow \mathbb{R}$ is defined as: $k(\mathbf{x}, \mathbf{x}') = \frac{1 - (\mathbf{x}^T \mathbf{x}')^p}{1 - (\mathbf{x}^T \mathbf{x}')$, where p is a non-negative integer. Which of the following is the corresponding feature mapping for this kernel when $D = 2$ and $p = 2$?

- (A) $\phi(\mathbf{x}) = [x_1, x_2, x_1 x_2]^T$
 (B) $\phi(\mathbf{x}) = [x_1, x_2, x_1^2, x_2^2]^T$
 (C) $\phi(\mathbf{x}) = [x_1, x_2, 1]^T$
 (D) $\phi(\mathbf{x}) = [x_1, x_2, x_1^2, x_2^2, x_1 x_2]^T$